INDEX TO VOLUME 27

Subject Index

Abortion

Bridging the gap: integrating family planning with abortion services in Turkey, see **Service Providers.**

Clandestine abortion in Latin America: a clinic profile, see **Latin America and the Caribbean.**

Court undercuts female feticide, Update, p. 111.

Effects of sex preference on contraceptive use, abortion and fertility in Matlab, Bangladesh, see **Asia**.

Incidence of induced abortion in Southern Ghana, see **Africa.**

Induced abortion in urban Nepal, Research Note, Thapa S and Padhye SM, p. 144.

Medical abortion at home, Update, p. 110.

Medical abortion regimen with reduced mifepristone dose, home misoprostol use is feasible and effective, Digest, p. 210.

Why Nigerian adolescents seek abortion rather than contraception: evidence from focus-group discussions, see **Africa.**

Africa

Acceptability of a vaginal microbicide among South African men, see **Contraceptive Research**.

Contraceptive use, intention to use and unmet need during the extended postpartum period, see **Contraceptive Practice.**

Despite symptoms, many Kenyans delay treatment for STDs, have unsafe sex, see **Sexually Transmitted Diseases**.

Douching is indirectly linked to HIV infection in female sex workers, see **Sexually Transmitted Diseases**.

Female circumcision in Sudan: future prospects and strategies for eradication, Islam MM and Uddin MM, p. 71.

Gender differences in adult perspectives on adolescent reproductive behaviors: evidence from Lomé, Togo, see **Sexual Behavior**.

Impact of a regional family planning service promotion initiative in Sub-Sarahan Africa: evidence from Cameroon, Babalola S, Vondrasek C, Brown J and Traoré R, p. 186.

In Kenya, the risk of poor birth outcomes is highest for a woman's first child, see **Pregnancy and Childbirth**.

In rural Gambia, men's marital and reproductive patterns differ sharply from those of women, Digest, p. 107.

In Sub-Saharan Africa, circumcised men are less likely than uncircumcised men to become infected with HIV, Digest, p. 102.

Incidence of induced abortion in southern Ghana, Ahiadeke C, p. 96.

Many women in rural Gambia have reproductive health problems, but few of them seek treatment, see **Reproductive Health**.

Medical abortion regimen with reduced mifepristone dose, home misoprostol use is feasible and effective, see **Abortion**.

Monitoring quality of care in family planning programs: a comparison of observations and client exit interviews, see **Program Evaluation and Funding.**

Obstetric complications common in West Africa, despite accessible care, see **Pregnancy and Childbirth.**

Promoting sexual responsibility among young people in Zimbabwe, Kim YM, Kols A, Nyakauru R, Marangwanda C and Chibatamoto P, p. 11.

Risk of HIV transmission is raised by high viral load, presence of genital ulcers, see **Sexually Transmitted Diseases**.

South Africans disagree with Mbeki, Update, p. 55.

Timing of first intercourse among Malian adolescents: implications for contraceptive use, Gueye M, Castle S and Konaté MK, p. 56.

Why do family planning providers restrict access to services? an examination in Ghana, Stanback J and Twum-Baah KA, p. 37.

Why Nigerian adolescents seek abortion rather than contraception: evidence from focus-group discussions, Otoide VA, Oronsaye F and Okonofua FE, p. 77.

Asia

Contraceptive use, intention to use and unmet need during the extended postpartum period, see **Contraceptive Practice.**

Consequences of the shift from domiciliary distribution to site-based family planning services in Bangladesh, see **Program Evaluation and Funding.**

Effects of sex preference on contraceptive use, abortion and fertility in Matlab, Bangladesh, Bairagi R, p. 137.

Indian women often select methods other than those their providers recommend, see **Service Providers.**

Induced abortion in urban Nepal, see **Abortion**.

Kinship networks and contraceptive choice in Nang Rong, Thailand, Godley J, p. 4.

Manila meeting focuses on equity, Update, p. 54.

Maternal zinc supplements in pregnancy lower risks linked to low birth weight, see **Maternal and Child Health**.

Medical abortion regimen with reduced mifepristone dose, home misoprostol use is feasible and effective, see **Abortion**.

Peer counselors' support is successful in promoting exclusive breastfeeding among Bangladeshi women, see **Breastfeeding.**

Peer training of nurses improves immunization coverage and practices, see **Service Providers**.

Persistence of a service delivery 'culture': findings from a qualitative study in Bangladesh, see **Service Providers**.

Prevention of unwanted births in India would result in replacement fertility, Digest, p. 104.

Thai men who patronize prostitutes place their wives at risk of HPV-associated cervical cancer, see **Reproductive Health**.

What influences contraceptive use among young women in urban squatter settlements of Karachi, Pakistan? see **Contraceptive Practice**.

Why are condoms used, and how many are need? estimates from Orissa, India, see **Barrier Methods**.

Barrier Methods

Condom use with risky partners, Update, p. 163.

Condoms reduce women's risk of herpes infection, but do not protect men, Digest, p. 213.

Female condom reuse, Update, p. 3.

Female condoms remain structurally sound after being washed and reused as many as seven times, Digest, p. 155.

Why are condoms used, and how many are needed? estimates from Orissa, India, Collumbien M, Das B and Campbell OMN, p. 171.

Why sex workers use condoms, Update, p. 3.

Breastfeeding

Breastfeeding and working mothers, Update, p. 54.

Characteristics of mother, child linked to postnatal HIV transmission risk, see **Maternal and Child Health.**

Mothers breastfeed, despite HIV, Update, p. 163.

Peer counselors' support is successful in promoting exclusive breastfeeding among Bangladeshi women, Digest, p. 51.

Contraceptive Practice

Condoms reduce women's risk of herpes infection, but do not protect men, see **Barrier Methods**.

Contraceptive dynamics in Guatemala: 1978–1998, Bertrand JT, Seiber E and Escudero G, p. 112.

Contraceptive use, intention to use and unmet need during the extended postpartum period, Ross JA and Winfrey WL, p. 20.

Effects of sex preference on contraceptive use, abortion and fertility in Matlab, Bangladesh, see **Asia**.

Female condoms remain structurally sound after being washed and reused as many as seven times, see **Barrier Methods**.

First pregnancy may be difficult to achieve after long-term use of an IUD, see **Intrauterine Devices**.

Impact of a regional family planning service

promotion initiative in Sub-Sarahan Africa: evidence from Cameroon, see **Africa.**

Indian women often select methods other than those their providers recommend, see **Service Providers.**

Kinship networks and contraceptive choice in Nang Rong, Thailand, see **Asia.**

Timing of first intercourse among Malian adolescents: implications for contraceptive use, see **Africa.**

What influences contraceptive use among young women in urban squatter settlements of Karachi, Pakistan? Fikree FF, Khan A, Kadir MM, Sajan F and Rahbar MH, p. 130.

Why are condoms used, and how many are needed? estimates from Orissa, India, see **Barrier Methods**.

Why Nigerian adolescents seek abortion rather than contraception: evidence from focus-group discussions, see **Africa.**

Contraceptive Research
Acceptability of a vaginal microbicide among South African men, Ramjee G, Gouws E, Andrews A, Myer L and Weber AE, p. 164.

Female condoms remain structurally sound after being washed and reused as many as seven times, see **Barrier Methods**.

Emergency Contraception
Most abortion in China avoidable, Update, p. 162.

Fertility: Planning and Effects
Contraceptive use, intention to use and unmet need during the extended postpartum period, see **Contraceptive Practice.**

Effects of sex preference on contraceptive use, abortion and fertility in Matlab, Bangladesh, see **Asia**.

First pregnancy may be difficult to achieve after long-term use of an IUD, see **Intrauterine Devices**.

Prevention of unwanted births in India would result in replacement fertility, see **Asia.**

Hormonal Implants and Injectables
Counseling leads to continuation, Update, p. 110.

Intrauterine Devices
First pregnancy may be difficult to achieve after long-term use of an IUD, Digest, p. 158.

IUD and recovery of fertility, Letter, Sivin I, Webb A and Stedman Y, p. 208.

STDs, not IUDs, cause infertility, Update, p. 163.

Latin America and the Caribbean
Clandestine abortion in Latin America: a clinic profile, Strickler J, Heimburger A and Rodriguez K, p. 34.

Contraceptive dynamics in Guatemala: 1978–1998, see **Contraceptive Practice**.

Contraceptive use, intention to use and unmet need during the extended postpartum period, see **Contraceptive Practice.**

Length of counseling sessions and the amount of relevant information exchanged: a study in Peruvian clinics, León FR, Monge R, Zumarán A, García I and Ríos A, p. 28; and Correction, p. 89.

Monitoring quality of care in family planning programs: a comparison of observations and client exit interviews, see **Program Evaluation and Funding.**

Perceptions of peer behavior predict whether Peruvian adolescents have had sex, see **Sexual Behavior**.

Risk of mother-child transmission of syphilis is elevated if the woman has clinical signs of disease, see **Pregnancy and Childbirth.**

Strategies for detection of sexually transmitted infection among family planning clients in Jamaica, see **Sexually Transmitted Diseases**.

Males
Acceptability of a vaginal microbicide among South African men, see **Contraceptive Research**.

Condoms reduce women's risk of herpes infection, but do not protect men, see **Barrier Methods**.

In rural Gambia, men's marital and reproductive patterns differ sharply from those of women, see **Africa.**

In Sub-Saharan Africa, circumcised men are less likely than uncircumcised men to become infected with HIV, see **Africa.**

Men who have many sexual partners before marriage are more likely to engage in extramarital intercourse, Digest, p. 48.

Thai men who patronize prostitutes place their wives at risk of HPV-associated cervical cancer, Digest, see **Reproductive Health**.

Why are condoms used, and how many are needed? estimates from Orissa, India, see **Barrier Methods**.

Maternal and Child Health
Characteristics of mother, child linked to postnatal HIV transmission risk, Digest, p. 103.

Female circumcision in Sudan: future prospects and strategies for eradication, see **Africa.**

Maternal zinc supplements in pregnancy lower risks linked to low birth weight, Digest, p. 213.

Mortality under five, Update, p. 3.

Nurturing the newly born, Update, p. 162.

Peer training of nurses improves immunization coverage and practices, see **Service Providers**.

Short anti-HIV treatment okay for infant if mother receives standard regimen, Digest, p. 159.

Middle East
Contraceptive use, intention to use and unmet need during the extended postpartum period, see **Contraceptive Practice.**

Population Policy and Research
Gagged again, Update, p. 2.

Picture this…, Letter, Rinehart W, p. 209.

Prevention of unwanted births in India would result in replacement fertility, see **Asia.**

Media, public opinion and population assistance: establishing the link, see **Public Opinion.**

Using photographs to strengthen family planning research, Viewpoint, Donaldson PJ, p. 148.

Pregnancy and Childbirth
Disparities in pregnancy risks, Update, p. 55.

Fewer prenatal visits can be fine, Update, p. 111.

First pregnancy may be difficult to achieve after long-term use of an IUD, see **Intrauterine Devices**.

For breech births, cesarean section poses fewer risks than vaginal delivery, Digest, p. 105.

HIV therapy options in pregnancy, Update, p. 54.

In Kenya, the risk of poor birth outcomes is highest for a woman's first child, Digest, p. 156.

Malaria risk after delivery, Update, p. 2.

Maternal zinc supplements in pregnancy lower risks linked to low birth weight, see **Maternal and Child Health**.

Obstetric complications common in West Africa, despite accessible care, Digest, p. 52.

Pregnant adolescents at risk, Update, p. 110.

Pregnant women experiencing job-related fatigue face elevated risk of premature membrane rupture, Digest, p. 160.

Risk of mother-child transmission of syphilis is elevated if the woman has clinical signs of disease, Digest, p. 106.

Unsafe pregnancy intervals, p. 2.

Program Evaluation and Funding
Bridging the gap: integrating family planning with abortion services in Turkey, see **Service Providers.**

Consequences of the shift from domiciliary distribution to site-based family planning services in Bangladesh, Routh S, Ashraf A, Stoeckel J and Barkat-e-Khuda, p. 82.

Family planning program effort index: 1999 cycle, Ross J and Stover J, p. 119.

Length of counseling sessions and the amount of relevant information exchanged: a study in Peruvian clinics, see **Latin America and the Caribbean.**

Monitoring quality of care in family planning programs: a comparison of observations and client exit interviews, Bessinger RE and Bertrand JT, p. 63.

Promoting sexual responsibility among young people in Zimbabwe, see **Africa**.

Personalized HIV counseling and testing show promise in reducing risk behaviors, see **Sexually Transmitted Diseases.**

Public Opinion
Acceptability of a vaginal microbicide among South African men, see **Contraceptive Research**.

Media, public opinion and population assistance: establishing the link, Viewpoint, Schindlmayr T, p. 42.

Reproductive Health

Douching is indirectly liked to HIV infection in female sex workers, see **Sexually Transmitted Diseases**.

Female circumcision in Sudan: future prospects and strategies for eradication, see **Africa.**

In rural Gambia, men's marital and reproductive patterns differ sharply from those of women, see **Africa.**

Many women in rural Gambia have reproductive health problems, but few of them seek treatment, Digest p. 214.

Risk factors for low-grade cervical abnormalities differ from those for human papillomavirus infection, Digest, p. 212.

Thai men who patronize prostitutes place their wives at risk of HPV-associated cervical cancer, Digest, p. 154.

Two-stage cervical cancer testing: more women can be screened, and fewer undergo unnecessary treatment, Digest, p. 47.

Research Methods

Monitoring quality of care in family planning programs: a comparison of observations and client exit interviews, see **Program Evaluation and Funding.**

Survey respondents like computers, Update, p. 54.

Service Providers

Bridging the gap: integrating family planning with abortion services in Turkey, Comment, Senlet P, Cagatay L, Ergin J and Mathis J, p. 90.

Busy obstetricians favor cesareans, Update, p. 55.

Can pharmacists treat STIs? Update, p. 162.

Clandestine abortion in Latin America: a clinic profile, see **Latin America and the Caribbean.**

Consequences of the shift from domiciliary distribution to site-based family planning services in Bangladesh, see **Program Evaluation and Funding.**

Contraceptive dynamics in Guatemala: 1978–1998, see **Contraceptive Practice**.

Gaps in HIV care in Central America, Update, p. 163.

Impact of a regional family planning service promotion initiative in Sub-Sarahan Africa: evidence from Cameroon, see **Africa**.

Indian women often select methods other than those their providers recommend, Digest, p. 50.

Length of counseling sessions and the amount of relevant information exchanged: a study in Peruvian clinics, see **Latin America and the Caribbean.**

Peer training of nurses improves immunization coverage and practices, Digest, p. 211.

Persistence of a service delivery 'culture': findings from a qualitative study in Bangladesh, Schuler SR, Bates LM and Islam MK, p. 194.

Provider perspective: human after all, Viewpoint, Shelton JD, p. 152.

Strategies for detection of sexually transmitted infection among family planning clients in Jamaica, see **Sexually Transmitted Diseases**.

Two-stage cervical cancer testing: more women can be screened, and fewer undergo unnecessary treatment, see **Reproductive Health**.

Why do family planning providers restrict access to services? an examination in Ghana, **see Africa.**

Sexual Behavior

Gender differences in adult perspectives on adolescent reproductive behaviors: evidence from Lomé, Togo, Speizer IS, Mullen SA and Amégee K, p. 178.

In rural Gambia, men's marital and reproductive patterns differ sharply from those of women, see **Africa.**

Men who have many sexual partners before marriage are more likely to engage in extramarital intercourse, see **Males.**

Perceptions of peer behavior predict whether Peruvian adolescents have had sex, Digest, p. 157.

Promoting sexual responsibility among young people in Zimbabwe, see **Africa**.

Risk factors for low-grade cervical abnormalities differ from those for human papillomavirus infection, see **Reproductive Health**.

Timing of first intercourse among Malian adolescents: implications for contraceptive use, see **Africa.**

Sexually Transmitted Diseases

AIDS update 2000, Update, p. 2.

Better HIV surveillance needed, Update, p. 162.

Condoms reduce women's risk of herpes infection, but do not protect men, see **Barrier Methods**.

Despite symptoms, many Kenyans delay treatment for STDs, have unsafe sex, Digest, p. 211.

Douching is indirectly liked to HIV infection in female sex workers, Digest, p. 215.

Herpes is common in Mexico, Update, p. 111.

In Sub-Saharan Africa, circumcised men are less likely than uncircumcised men to become infected with HIV, see **Africa.**

Novel HIV prevention, Update, p. 3.

Personalized HIV counseling and testing show promise in reducing risk behaviors, Digest, p. 49.

Risk factors for low-grade cervical abnormalities differ from those for human papillomavirus infection, see **Reproductive Health**.

Risk of HIV transmission is raised by high viral load, presence of genital ulcers, Digest, p. 156.

STDs drive rise in circumcision, Update, p. 162.

Strategies for detection of sexually transmitted infection among family planning clients in Jamaica, Ward E, Spruyt A, Fox L, Johnson L, Wong E, Behets F, Figueroa JP and Morris J, p. 201.

Treating partners of STD patients, Update, p. 55.

UN session on HIV and AIDS, Update, p. 111.

Status of Women

Costs of gender inequality, p. 3.

Female circumcision in Sudan: future prospects and strategies for eradication, see **Africa.**

Girls are worth the investment, Update, p. 110.

Manila meeting focuses on equity, see **Asia**.

Persistence of a service delivery 'culture': findings from a qualitative study in Bangladesh, see **Service Providers**.

What influences contraceptive use among young women in urban squatter settlements of Karachi, Pakistan? see **Contraceptive Practice**.

Teenagers

Gender differences in adult perspectives on adolescent reproductive behaviors: evidence from Lomé, Togo, see **Sexual Behavior**.

Perceptions of peer behavior predict whether Peruvian adolescents have had sex, see **Sexual Behavior**.

Promoting sexual responsibility among young people in Zimbabwe, see **Africa**.

Timing of first intercourse among Malian adolescents: implications for contraceptive use, see **Africa.**

Why Nigerian adolescents seek abortion rather than contraception: evidence from focus-group discussions, see **Africa.**

Author Index

Ahiadeke C, Incidence of induced abortion in southern Ghana, p. 96.

Amégee K, see Speizer IS.

Andrews A, see Ramjee G.

Ashraf A, see Routh S.

Babalola S, Vondrasek C, Brown J and Traoré R, The impact of a regional family planning service promotion initiative in Sub-Saharan Africa: evidence from Cameroon, p. 186.

Barkat-e-Khudu, see Routh S.

Bairagi R, Effects of sex preference on contraceptive use, abortion and fertility in Matlab, Bangladesh, p. 137.

Bates LM, see Schuler SR.

Behets F, see Ward E.

Bertrand JT, Seiber E and Escudero G, Contraceptive dynamics in Guatemala:1978–1998, p. 112.

Bertrand JT, see Bessinger RE.

Bessinger RE and Bertrand JT, Monitoring quality of care in family planning programs: a comparison of observations and client exit interviews, p. 63.

Brown J, see Babalola S.

Cagatay L, see Senlet P.

Campbell OMR, see Collumbien M.

Castle S, see Gueye M.

Chibatamoto P, see Kim YM.

Collumbien M, Das B and Campbell OMR, Why are condoms used, and how many are needed? estimates from Orissa, India, p. 171.

Das B, see Collumbien M.

Donaldson PJ, Using photographs to strengthen family planning research, Viewpoint, p. 148.

Ergin J, see Senlet P.

Escudero G, see Bertrand JT.

Figueroa JP, see Ward E.

Fikree FF, Khan A, Kadir MM, Sajan F and Rahbar MH, What influences contraceptive use among young women in urban squatter settlements of Karachi, Pakistan? p. 130.

Fox L, see Ward E.

García I, see León FR.

Godley J, Kinship networks and contraceptive choice in Nang Rong, Thailand, p. 4.

Gouws E, see Ramjee G.

Gueye M, Castle S and Konaté MK, Timing of first intercourse among Malian adolescents: implications for contraceptive use, p. 56.

Heimburger A, see Strickler J.

Islam MK, see Schuler SR.

Islam MM and Uddin MM, Female circumcision in Sudan: future prospects and strategies for eradication, p. 71.

Johnson L, see Ward E.

Kadir MM, see Fikree FF.

Khan A, see Fikree FF.

Kim YM, Kols A, Nyakauru R, Marangwanda C and Chibatamoto P, Promoting sexual responsibility among young people in Zimbabwe, p. 11.

Kols A, see Kim YM.

Konaté MK, see Gueye M.

León FR, Monge R, Zumarán A, García I and Ríos A, Length of counseling sessions and the amount of relevant information exchanged: a study in Peruvian clinics, p. 28.

Marangwanda C, see Kim YM.

Mathis J, see Senlet P.

Monge R, see León FR.

Morris J, see Ward E.

Mullen SA, see Speizer IS.

Myer L, see Ramjee G.

Nyakauru R, see Kim YM.

Okonofua FE, see Otoide VO.

Oronsaye F, see Otoide VO.

Otoide VO, Oronsaye F and Okonofua FE, Why Nigerian adolescents seek abortion rather than contraception: evidence from focus-group discussions, p. 77.

Padhye SM, see Thapa S.

Rahbar MH, see Fikree FF.

Ramjee G, Gouws E, Andrews A, Myer L and Weber AE, The acceptability of a vaginal microbicide among South African men, p. 164.

Rinehart W, Picture this…., Letter, p. 209.

Ríos A, see León FR.

Rodriguez K, see Strickler J.

Ross JA and Winfrey WL, Contraceptive use, intention to use and unmet need during the extended postpartum period, p. 20.

Ross J and Stover J, The family planning program effort index: 1999 cycle, p. 119.

Routh S, Ashraf A, Stoeckel J and Barkat-e-Khuda, Consequences of the shift from domiciliary distribution to site-based family planning services in Bangladesh, p. 82.

Sajan F, see Fikree FF.

Schindlmayr T, The media, public opinion and population assistance: establishing the link, Viewpoint, p. 42.

Schuler SR, Bates LM and Islam MK, The persistence of a service delivery 'culture': findings from a qualitative study in Bangladesh, p. 194.

Seiber E, see Bertrand JT.

Senlet P, Cagatay L, Ergin J and Mathis J, Bridging the gap: integrating family planning with abortion services in Turkey, Comment, p. 90.

Shelton JD, The provider perspective: human after all, Viewpoint, p. 152.

Sivin I, Webb A and Stedman Y, The IUD and recovery of fertility, Letter, p. 208.

Speizer IS, Mullen SA and Amégee K, Gender differences in adult perspectives on adolescentreproductive behaviors: evidence from Lomé, Togo, p. 178.

Spruyt A, see Ward E.

Stanback J and Twum-Baah KA, Why do family planning providers restrict access to services? an examination in Ghana, Research Note, p. 37.

Stedman Y, see Sivin I.

Stoeckel J, see Routh S.

Stover J, see Ross J.

Strickler J, Heimburger A and Rodriguęz K, Clandestine abortion in Latin America: a clinic profile, Research Note, p. 34.

Thapa S and Padhye SM, Induced abortion in urban Nepal, Research Note, p. 144.

Traoré R, see Babalola S.

Twum-Baah KA, see Stanback J.

Uddin MM, see Islam MM.

Vondrasek C, see Babalola S.

Ward E, Spruyt A, Fox L, Johnson L, Wong E, Behets F, Figueroa JP and Morris J, Strategies for detection of sexually transmitted infection among family planning clients in Jamaica, p. 201.

Webb A, see Sivin I.

Weber AE, see Ramjee G.

Winfrey WL, see Ross JA.

Wong E, see Ward E.

Zumarán A, see León FR.

Statement of Ownership, Management and Circulation

Title of publication: *International Family Planning Perspectives.* **Publication no.:** 404170. **Date of filing:** Oct. 1, 2001. **Frequency of issue:** Quarterly. **No. of issues published annually:** Four. **Annual subscription price:** Individuals, $36.00; Institutions, $46.00. **Complete mailing address of known office of publication:** 120 Wall Street, New York, NY 10005. **Complete mailing address of the headquarters or general business offices of the publisher:** Same as above. **Publisher:** The Alan Guttmacher Institute, same address as above. **Editor:** Patricia Donovan, same address as above. **Owner:** The Alan Guttmacher Institute. **Known bondholders, mortgagees, and other security holders owning or holding 1 percent or more of total amount of bonds, mortgages or other securities:** None.

The purpose, function, and nonprofit status of this organization and the exempt status for federal income tax purposes have not changed during preceding 12 months.

Extent and nature of circulation	Avg. no. copies each issue during preceding 12 mos.	Actual no. copies of single issue published nearest to filing date
Total no. copies	31,013	33,200
Paid and/or requested circulation		
Sales through dealers and carriers, street vendors, and counter sales	0	0
Mail subscription (paid and/or requested)	29,914	32,036
Total paid and/or requested circulation	29,914	32,036
Free distribution by mail, carrier or other means; samples, complimentary and other free copies	899	964
Total distribution	30,813	33,000
Copies not distributed		
Office use, left over, unaccounted for, spoiled after printing	200	200
Return from news agents	0	0
Total	31,013	33,200

I certify that the statements made by me above are correct and complete.

Patricia Donovan, Editor